


Psychometric properties of the Minnesota Living with Chronic Heart Failure Questionnaire in a Colombian population

Propiedades psicométricas del “Minnesota Living with Chronic Heart Failure Questionnaire” en una Población Colombiana

Lyda Z. Rojas^{1,2}, Juliana A. Hernández¹, Silvia J. Trujillo¹, Zayne M. Roa¹, Adriana M. Jurado¹,
Luis E. Echeverría¹, Taulant Muka², Oscar H. Franco^{2,3}

Suggested citation: Rojas LZ, Hernández JA, Trujillo SJ, Roa ZM, Jurado AM, Echeverría LE, et al. Psychometric properties of the Minnesota Living with Chronic Heart Failure Questionnaire in a Colombian population. Rev Univ Ind Santander Salud. 2019;51(1): 11-22. doi: 10.18273/revsal.v51n1-2019002 

Abstract

Introduction: Although the Minnesota Living with Heart Failure Questionnaire (MLHF-Q) is one of the most widely used tools to assess Health-Related Quality of Life (HRQoL) in patients with chronic heart failure (CHF), it has not been validated in Latin American Spanish-speaking populations. **Objective:** We evaluated internal consistency and construct validity of the MLHF-Q in patients with CHF from Colombia. **Methods:** The Spanish version of the MLHF-Q was given to 200 patients. Cronbach's alpha was used to evaluate internal consistency. Confirmatory factorial Principal Component Analysis (PCA) and Rasch analysis were used to evaluate construct validity. The discriminative capacity was measured using the Mann-Whitney U test. **Results:** Median age was 64 years, 63% of the patients included in the study were men, and 79.5% had a left ventricular ejection fraction (LVEF) \leq 45%. The median of the total score of HRQoL was 40 points (Q1=20; Q3=55), physical dimension 11 points (Q1=4; Q3=23) and emotional dimension 7 points (Q1=3; Q3=13). Global internal consistency of MLHF-Q was 0.91 (95% CI 0.89 - 0.93). In the PCA, the three dimensions explained 47.7% and 54.0% in Rasch analysis, in which five items presented misfit. Worse HRQoL was observed among women than men in the emotional dimension ($p=0.047$). Discriminative capacity for the overall score of the MLHF-Q and their subscales was observed in age and New York Heart Association (NYHA) functional class ($p<0.05$). **Conclusions:** Our findings confirmed the three-factor structure of the MLHF-Q, and satisfactory level for internal consistency. Additionally, these results suggest that the questionnaire adequately reflects the severity of the disease. However, further studies are required to validate these findings in Colombian population and to evaluate the sensitivity to change of the MLHF-Q in longitudinal designs.

Keywords: Heart failure, quality of life, validation studies.

1. Fundación Cardiovascular de Colombia, Floridablanca, Santander, Colombia.

2. Erasmus University Medical Center, Rotterdam, The Netherlands.

3. University of Bern, Bern, Switzerland.

Correspondence: Lyda Z. Rojas. Address: Calle 155A No 23-58 Urbanización El Bosque. Box 681001 Floridablanca, Colombia. Phone number: 6399292 Ext. 344. Email: lydarojas@fcv.org

Resumen

Introducción: El Minnesota Living with Heart Failure Questionnaire (MLHF-Q) es uno de los instrumentos más utilizados para medir la Calidad de Vida Relacionada con la Salud (CVRS) en pacientes con Falla cardíaca Crónica (FCC); sin embargo, éste no ha sido validado en poblaciones latinoamericanas hispanohablantes. **Objetivo:** Evaluar la consistencia interna y validez de constructo del MLHF-Q en pacientes con FCC de Colombia. **Métodos:** La versión en español del MLHF-Q fue diligenciada por 200 pacientes. La consistencia interna se evaluó con el Alpha de Cronbach. La validez de constructo fue examinada por dos métodos: Análisis de Componentes Principales (ACP) confirmatorio y análisis Rasch. Se evaluó la capacidad discriminativa del instrumento con la prueba U-Mann-Whitney. **Resultados:** Mediana de edad de 64 años, 63% hombres y el 79.5% de los participantes tenían fracción de eyección del ventrículo izquierdo (FEVI) $\leq 45\%$. La mediana del puntaje total de CVRS fue 40 puntos (Q1=20; Q3=55), dimensión física 11 puntos (Q1=4; Q3=23) y dimensión emocional 7 puntos (Q1=3; Q3=13). La consistencia interna fue 0.91 (IC 95% 0.89 - 0.93). En el ACP, las tres dimensiones explicaron el 54.0% y 47.7% en el análisis Rasch, en éste último cinco ítems presentaron desajuste. Se observó peor CVRS en mujeres que en hombres en la dimensión emocional ($p=0.047$) y se evidenció capacidad discriminativa de las subescalas y el puntaje total del MLHF-Q en la edad y la clase funcional New York Heart Association (NYHA) ($p<0.05$). **Conclusión:** Nuestros hallazgos confirmaron la estructura de tres factores del MLHF-Q y un nivel satisfactorio para la consistencia interna. Adicionalmente, estos resultados sugieren que el cuestionario refleja adecuadamente la gravedad de la enfermedad. Sin embargo, se requieren estudios adicionales en población colombiana para validar estos hallazgos y evaluar la sensibilidad al cambio del MLHF-Q en diseños longitudinales.

Palabras clave: Insuficiencia cardíaca, calidad de vida, estudios de validación.

Introduction

Chronic Heart Failure (CHF) prevalence is over 5.8 million in the USA, and over 23 million worldwide¹. In Latin America, estimated Heart Failure (HF) prevalence is 1% (95% CI, 0.1 to 2.7%) with 199 per 100,000 person-year². The hospital readmission rates of patients with HF are 33%, 28%, 31%, and 35% at 3, 6, 12, and 24 to 60 months of follow-up (median duration of hospitalization of 7.0 days), with 1-year mortality rate of 24.5% (95% CI, 19.4 to 30.0%) and in-hospital mortality of 11.7% (95% CI, 10.4 to 13.0%)². Although mortality has decreased in CHF, the estimated survival rate is 50% and 10% at five and ten years after diagnosis³.

The burden of the disease in CHF involves several limitations in patients carrying out daily life activities, and affects health-related quality of life (HRQoL) more severely than other chronic diseases⁴. Several studies have shown that HRQoL of patients with HF is worse than the general population, or patients with other chronic diseases^{4,5}. Furthermore, the decline in quality of life of HF patients is not temporary, but rather progressive over time⁶. Nevertheless, measuring HRQoL in HF remains a challenge, and despite the existence of several instruments (generic and disease-specific) for assessing HRQoL, no consensus has been achieved on which instrument would be most suitable⁷.

The Minnesota Living with Heart Failure Questionnaire (MLHF-Q) is a disease-specific instrument, consisting of 21 items addressing a wide range of HRQoL and it is the most frequently used internationally. Since 1987, the MLHF-Q has been translated into more than 30 languages, including Spanish⁸⁻¹³ and it is used as an outcome measure in multiple clinical trials showing the best psychometric properties as to validity, reliability and sensitivity to change¹⁴⁻¹⁷.

Even though Spanish is spoken by 95% of Latin America's population, Brazil, where Portuguese is spoken, is the only regional country where the MLHF-Q has been validated¹⁸, while Colombia has no data available on the evaluation of the reliability and validity of the MLHF-Q. Therefore, we aimed to evaluate the internal consistency and construct validity of the MLHF-Q in patients with CHF in Colombia.

Methods

Study population

A cross-sectional study was conducted between February and October 2015 in the Heart Failure and Heart Transplant Clinic of Cardiovascular Foundation in Floridablanca city, Santander-Colombia. We included patients if they (i) were 18 years old or older and (ii) had a confirmed HF medical diagnosis by a cardiologist. Patients with mental sphere alterations or communication

limitations were excluded. All patients gave written informed consent and the Research Ethic Committee of the institution approved the research protocol.

In calculating our sample size, care was taken to comply with the 10-patient-per-analyzed-item criterion considered adequate for factorial analysis¹⁹. The sample was selected in a non-probabilistic way; all patients were invited to participate consecutively by a previously trained nurse who conducted the interviews upon medical control appointments.

Clinical screening

HRQoL was measured with the MLHF-Q¹¹, a specific self-report instrument for CHF patients. HRQoL questionnaire is made up of 21 items graded by the patient using a 6-point Likert-type scale ranging from 0 (no impairment) to 5 (very much impairment). The MLHF-Q groups the items in three dimensions: physical (8 items), emotional (5 items), and the overall score for HRQoL (21 items). Eight separate items, which do not assess a single construct or dimension of HRQoL, measure social and economic impairments of patients with HF and contribute to the overall score. The total score has a range between 0 and 105 points, the physical dimension (between 0 and 40), the emotional dimension (0 and 25) and the separate items on the socio-economic impairments (0 and 40). High scores on the MLHF-Q scale indicate impaired HRQoL. The MLHF-Q has a global internal consistency measured by Cronbach's alpha of 0.94 (95% CI, 0.91 to 0.95) and general intraclass correlation coefficient of 0.84, characteristics that make it suitable for use¹⁷.

Statistical analysis

Continuous variables are reported as median and quartiles (Q) unless stated otherwise, and categorical variables are presented as percentages. Internal consistency was evaluated through Cronbach's alpha coefficient²⁰. Kaiser-Meyer-Olkin's (KMO) index and Bartlett's test of sphericity were estimated to establish the pertinence of factorial analysis. KMO ≥ 0.7 was considered acceptable^{21,22}.

To evaluate construct validity of the questionnaire, two different approaches were used: first, the structure of the model originally proposed by Rector and Cohn⁸ was examined by means of confirmatory factorial principal component analysis (PCA). Dimensional structure was identified through varimax-type octagonal rotation, factor loading, and those ≥ 0.4 were considered

acceptable^{13,23}. Second, polytomous Rasch rating scale model was used to assess each specific questionnaire dimension according to the factorial structure proposed by literature²⁴. Thus, the first step was to evaluate the functioning of rating scale categories. A clearly progressive level of difficulty across item categories was expected as a criterion of adequate function. We also examined the standardized (ZSTD) fit statistics of persons for whom a score between ± 3 was expected.

For dimensionality evaluation, which is a fundamental requirement for construct validity, we applied the following criteria: (i) mean square information-weighted statistic (infit) and the outlier-sensitive statistic (outfit), with values between 0.7 and 1.3 indicate a good fit (ii) PCA of the residuals²⁵. Unidimensionality was violated if, besides the first factor, other factors had eigenvalues >3 , and the local dependency was assessed through the item residual correlations where values >0.5 may indicate that the response to one item may be determined by another. To detect the presence of differential item functioning (DIF), which occurs when groups within the sample respond differently from an individual item; we compared distinct levels of the trait by sex and age group (≤ 65 vs. >65 years). A Welch's t statistically significant ($p < 0.05$), and a difficulty difference ≥ 0.5 logits were considered evidence of uniform DIF.

Finally, discriminative capacity of the questionnaire was assessed by its ability to differentiate among subgroups of patients with different levels of CHF severity, taking into account the following hypothesis: women, higher age, superior New York Heart Association (NYHA) functional class and Left Ventricular Ejection Fraction (LVEF) under 45% will have higher scores of the MLHF-Q, by using the Mann-Whitney U test. All statistical tests were two-sided and a p -value < 0.05 was considered significant. Data were analyzed using Stata Statistical Software, version 14 and Winsteps 3.80.0.

Results

Characteristics of the study population

The proportion of missing data was 0%. During recruitment period, two hundred CHF patients fulfilled the selection criteria, agreed to participate and completed the questionnaire. Median age of participants was 64 (Q1=53; Q3=73) years old, 63.0 % were men, 79.5% had a LVEF $\leq 45\%$, and 24.0% subjects were in NYHA functional class III-IV. Sociodemographic and clinical characteristics of the study population are shown in **Table 1**.

Table 1. Sociodemographic and clinical characteristics of the study population (n=200).

Variable	n (%)
Age (years) ^a	64 (53-73)
Sex	
Men	126 (63.0)
Women	74 (37.0)
Socioeconomic status	
Low	124 (62.0)
Medium	64 (32.0)
High	12 (6.0)
Education status	
Less than high school	132 (66.0)
High school	40 (20.0)
Greater than high school	28 (14.0)
Marital status	
Married/living in a free union	134 (67.0)
Widowers	27 (13.5)
Single	24 (12.0)
Divorced	15 (7.5)
Employment status	
Unemployed	71 (35.5)
Employees	50 (25.0)
Pensioner	43 (21.5)
Housewives	22 (11.0)
Students	14 (7.0)
Etiology of Chronic heart failure	
Ischemic	60 (30.0)
Chagas disease	40 (20.0)
Idiopathic	29 (14.5)
Multifactorial	27 (13.5)
Hipertensive	14 (7.0)
Valvular	12 (6.0)
Other (congenital, viral, peripartum and others)	18 (9.0)
LVEF (%)	
≤45	159 (79.5)
>45	41 (20.5)
Functional class NYHA	
I-II	152 (76.0)
III-IV	48 (24.0)
Charlson comorbidity index	
1	101 (50.5)
2	58 (29.0)
3	27 (13.5)
>3	14 (7.0)
Medications	
Beta-Blocker	194 (97.0)
ACE inhibitor	68 (34.0)
Angiotensin receptor blocker	110 (55.0)
Aldosterone antagonist	165 (82.5)
Diuretic	126 (63.0)
Digoxin	58 (29.0)

Source: authors. ^aMedian (first and third quartile). LVEF=Left ventricular ejection fraction; NYHA=New York Heart Association; ACE=Angiotensin-converting enzyme.

Psychometric analysis

Internal reliability

Cronbach's alphas coefficients ranged from 0.73 (social dimension) to 0.91 (physical dimension and total score) in the MLHF-Q, indicating satisfactory level for internal consistency. Descriptive analysis and internal consistency of the MLHF-Q are shown in **Table 2 and Supplementary Material Table S1**.

Table 2. Descriptive analysis and internal consistency of the Minnesota Living with Heart Failure Questionnaire (n=200).

Quality of life	Number of items	Median (Q1-Q3) ^a	Range	Cronbach's	95% CI
Physical dimension	8	11 (4-23)	0-40	0.91	0.89-0.93
Emotional dimension	5	7 (3-13)	0-24	0.80	0.76-0.85
Social	8	16 (10-22)	0-32	0.73	0.68-0.78
Total score	21	40 (20-55)	0-95	0.91	0.90-0.93

Source: authors. ^aQ1-Q3=First and third quartile.

Construct validity

The KMO statistic was 0.90, indicating sampling adequacy (**Supplementary Material Table S2**) and Bartlett's test of sphericity was statistically significant ($\chi^2(210)=2126.20$; $p=0.000$), suggesting that data were appropriate to be subjected to a factorial analysis²². All items in the first factor were associated to signs and symptoms of HF; this factor was identified such as physical dimension. The second factor, included four items of five items from the original questionnaire, and they were related to the patient's psychological response to disease; this factor was recognized as the emotional dimension. Finally, three items in the third factor were correlated to the patient's social relationships, thus this factor was named the social dimension. Then, confirmatory factorial PCA of three factors explained 54.03% of total variation in the study population, of which 30.6% was explained by the first factor, 15.8% the second factor and 7.6% the third dimension. Eigenvalue was 6.43 for the physical, 3.31 for the emotional, and 1.59 for the social dimension (**Supplementary Material Table S3**).

Table 3 shows factorial analysis, five of the 21 items demonstrated factor loadings between 0.4 and 0.6; four between 0.6 and 0.7; six between 0.7 and 0.8; four items had factor loadings >0.8 and, two items (14 and 16) did not adequately load (loading <0.4). Additionally, in **Table 3** we can distinguish physical,

Psychometric properties of the Minnesota Living with Chronic Heart Failure Questionnaire in a Colombian population

emotional and social dimensions. Also, items 1, 8, 9, 10 (social) and 20 (emotional) have been reclassified in physical dimension. Conversely, emotional dimension

is preserved almost entirely, except for item 20. Finally, item 8 could belong to both, the physical and social dimension.

Table 3. Factorial principal component analysis with orthogonal varimax rotation (loadings) of the Minnesota Living with Heart Failure Questionnaire.

Item	Factor1 (Physical)	Factor2 (Emotional)	Factor3 (Social)	Uniqueness
1. Swelling in your ankles, legs	0.5120 ^a			0.6223
2. Resting during day	0.7834 ^b			0.3565
3. Walking or climbing stairs difficult	0.8388 ^b			0.2816
4. Working around house difficult	0.8264 ^b			0.2747
5. Away from home difficult	0.7965 ^b			0.3158
6. Sleeping difficult	0.5547 ^b			0.6568
7. Relating to or doing things with friends or family difficult	0.6013 ^b			0.5491
8. Working to earn a living difficult	0.7058 ^a		0.4391 ^a	0.2883
9. Recreational activities difficult	0.6374 ^a			0.4775
10. Sexual activities difficult	0.4471 ^a			0.7653
11. Eating less foods I like			0.6183 ^a	0.5349
12. Shortness of breath	0.6765 ^b			0.4462
13. Fatigue	0.8216 ^b			0.2710
14. Hospitalization	----	----	----	0.7383
15. Medical costs			0.7819 ^a	0.3630
16. Side effects from medications	----	----	----	0.8387
17. Feeling burden to family or friends		0.7972 ^c		0.3314
18. Feeling loss of self-control		0.8719 ^c		0.2252
19. Being worried		0.7854 ^c		0.3146
20. Difficulty concentrating or remembering	0.4947 ^c			0.6825
21. Being depressed		0.7746 ^c		0.3204

Source: authors. ^aSingle items used in the construction of the overall score (social dimension); ^bPhysical dimension; ^cEmotional dimension.

Regarding Rasch analysis of the total score, the average measures of the rating scale of the MLHF-Q were ordered, progressing from -0.84 logits for rating scale category zero (no impairment) to 0.27 logits for rating scale category of five (very much impairment); disordered thresholds (response categories not working logically) were corrected by combining adjacent categories (**Supplementary Material Figure S1**); the result was a 3-point scale that met the criteria for rating scale. Eight persons had a ZSTD exceeding the value expected and were excluded from the analysis.

The person separation was 2.56 and reliability 0.87; for items these statistics were 5.00 and 0.96, respectively. Items 15, 16, 20, 14 and 10 showed fit statistics (Outfit, Infit) out of the established range for the analysis; statistics are shown in **Table 4**. In the PCA of residuals, 21 items and 192 persons explained the 42.0%. In

the first contrast, we observed 3.3 eigenvalues with residuals correlations higher than 0.50. Items 1, 15, 16 and 17 had a difference in difficulty ≥ 0.5 logits by groups of age with Welch p values under 0.05. There was no evidence of uniform DIF by sex groups. After removing the five misfit items, the overall fit of the data improved, with 47.7% of raw variance explained and only two items (6 and 11) maintaining their fit statistics above the range.

In the analysis of physical, emotional and social dimensions, disorders of the rating scale were not observed, in fact, all analyses were made with the original MLHF-Q codification. Social dimension's items explained 44.8% of the variance and had 1.7 eigenvalues in the first contrast, the residual did not present any important correlation. Item 8 (working to earn a living difficult) of social dimension had a slightly

lower value of the range. The eight items of physical dimension presented 2 eigenvalues in the first contrast and explained 57.4% of the raw variance; two of the

items were above and one item was below expected range as shown in **Table 4**.

Table 4. Severity levels, standard errors, and goodness-of-fit indexes of the 21 MLHF-Q items and by dimensions using Rasch analysis (n=192).

Items	Description	(logit) ^a	SE	MNSQ	
				Infit	Outfit ^b
Total					
15	Medical costs	-1.04	0.11	1.57	1.91
16	Side effects from medications	0.30	0.11	1.28	1.59
20	Difficulty concentrating or remembering	-0.38	0.11	1.13	1.56
14	Hospitalization	1.07	0.13	1.48	1.41
10	Sexual activities difficult	-0.34	0.11	1.35	1.33
6	Sleeping difficult	0.09	0.11	1.13	1.20
11	Eat less food you like	-1.58	0.12	1.17	1.10
9	Recreational activities difficult	-0.40	0.11	1.10	1.09
18	Feeling a loss of self-control	0.49	0.12	1.10	1.07
17	Feeling burden to family or friends	0.63	0.12	1.08	0.96
1	Swelling in your ankles, legs	0.90	0.13	1.01	0.98
19	Being worried	-0.09	0.11	1.00	0.99
7	Relating to or doing things with friends or family difficult	0.60	0.12	0.92	0.83
21	Being depressed	0.40	0.12	0.89	0.88
8	Working to earn a living difficult	-0.65	0.11	0.81	0.73
3	Walking or climbing stairs difficult	-0.43	0.11	0.77	0.72
5	Being away from home difficult	0.14	0.11	0.77	0.66
12	Shortness of breath	0.53	0.12	0.73	0.76
4	Working around house difficult	-0.02	0.11	0.73	0.65
2	Resting during day	0.04	0.11	0.60	0.57
13	Fatigue	-0.25	0.11	0.57	0.56
Physical					
6	Sleeping difficult	-0.04	0.06	1.70	1.88
7	Relating to or doing things with friends or family difficult	0.39	0.07	1.43	1.55
12	Shortness of breath	0.33	0.07	1.07	1.26
5	Being away from home difficult	0.03	0.07	0.95	0.83
4	Working around house difficult	-0.05	0.06	0.85	0.81
3	Walking or climbing stairs difficult	-0.36	0.06	0.80	0.75
2	Resting during day	-0.06	0.06	0.78	0.75
13	Fatigue	-0.24	0.06	0.60	0.65
Emotional					
20	Difficulty concentrating or remembering	-0.43	0.06	1.72	2.30
17	Feeling burden to family or friends	0.30	0.07	0.93	0.86
19	Being worried	-0.19	0.06	0.80	0.77
21	Being depressed	0.09	0.07	0.74	0.64
18	Feeling a loss of self-control	0.23	0.07	0.70	0.62
Social					
10	Sexual activities difficult	-0.08	0.05	1.25	1.29
16	Side effects from medications	0.24	0.05	1.10	1.28
14	Hospitalization	0.51	0.06	1.18	1.14
15	Medical costs	-0.33	0.05	1.07	1.17
9	Recreational activities difficult	-0.08	0.05	0.98	0.93
11	Eat less food you like	-0.55	0.05	0.86	0.90
1	Swelling in your ankles, legs	0.47	0.05	0.87	0.82
8	Working to earn a living difficult	-0.20	0.05	0.68	0.63

Source: authors. ^aδ level of severity (higher values indicate higher severity). ^bExpected range 0.7 - 1.3. **MLHF-Q**=Minnesota Living with Heart Failure Questionnaire; **SE**=Standard error; **MNSQ**=Mean square fit statistic; **Infit**=Inlier-sensitive fit; **Outfit**=Outlier-sensitive fit.

In the emotional dimension, one item presented a severe misfit **Table 4**, the variance explained by these items was 54.0% and had 1.7 eigenvalues in the first contrast; correlation of -0.51 between items 19 and 20 was found. Eliminating item 20 and analyzing the remaining four items, all statistics were into the expected values (**Supplementary Material Table S4**) and the variance explained improved (**Supplementary Material Table S5**). It was not detected uniform DIF by sex or age group in any dimension. Wright maps are presented for each dimension evaluated (**Supplementary Material Figure S2**).

Contrast validity

Discriminative capacity of the MLHF-Q subscales and for the overall score was observed in age and NYHA functional class ($p < 0.05$). Worse HRQoL was observed among women than men in the emotional dimension ($p = 0.047$). Although higher HRQoL impairment was evident in LVEF $\leq 45\%$ compared with LVEF $> 45\%$ patients, it was not statistically significant (**Supplementary Material Table S6**).

Discussion

To the best of our knowledge, the present work is the first study that has assessed the psychometric properties of the MLHF-Q in a Spanish-speaking population of Latin America. We have evaluated the internal consistency, construct validity through the two methods (PCA and Rasch analysis), and the discriminative capacity of the MLHF-Q in outpatients with CHF in Colombia.

Interpretation of findings

The reliability of subscales and overall MLHF-Q showed a Cronbach's alpha acceptable to excellent, with coefficients similar to those in other populations: Australia, France, Hungary, Yugoslavia (physical dimension $\alpha = 0.91$); Hungary, Poland, Sweden (emotional dimension $\alpha = 0.80$); Israel, Italy (social dimension $\alpha = 0.73$), and Denmark, Spain, Yugoslavia (total score $\alpha = 0.91$)^{23,26}.

Regarding construct validity, we found that the three factors explain 47.7% and 54.03% of the overall score in the Rasch analysis and PCA, respectively, with similar results previously reported by another study²⁷, while in other studies the variance explained by these three factors has been higher (64.1 to 72%)^{9,10,19,28}. Also, we found the following similarities with other authors; Heo, et al.²⁷ evidenced that items (1 and 9) were loaded on physical dimension and items (14 and 16) presented

loading < 0.4 . Ho, et al.¹⁹ showed that item 1 was loaded on physical dimension. Finally, Moon, et al.²⁸ found that items (1, 9 and 10) were loaded on physical dimension.

Item 1 (Swelling in your ankles, legs) is part of the social dimension (another dimension) from original version; however, it has been reported that up to two thirds of patients admitted with acute HF presented hypervolemia signs such as jugular venous distension and peripheral edema, typical physiopathological manifestations of HF²⁹⁻³⁰, which support its correlation to the physical dimension. On the other hand, determine the most plausible dimension for item 10 (sexual activities difficult) is complicated, due to the multifactorial explanation (psychological, emotional, physical and medical) of HF patients' sexual activity³¹. Also, possible explanations for the differences found in the factor structure, variance explained, and eigenvalues with other authors could be sample size, culture, demographics and clinical characteristics, among others^{10,17}.

The MLHF-Q is interpreted by its total score, which results of averaging the score of all 21 items. However, this assumes that the total score is unidimensional. Nevertheless, Rasch analysis for the total score did not find evidence of unidimensional functioning. Moreover, it demonstrated misfitting of five items (10, 14, 15, 16, 20), and therefore confirming the existence of some problematic items in the composition of the total score. Elimination of items has been reported as a solution^{10,27}. Exclusion of problematic items in our study improved the general fit to the Rasch model.

Similar findings have been reported by Munyombwe, et al.¹⁰ who found that several items (7, 8, 10, 14, 16) presented misfit. Also, Bilbao, et al.²⁴ reported two misfitting items (1 and 10). Considering that misfitting items have been identified in a third factor presenting the social dimension, as also shown in the current study, several authors have suggested to add a third factor to the total score^{9,10,19,23,28}. However, it remains a challenge to reach a consensus on which of the different social factors proposed is the most appropriate and has the best psychometric properties, and therefore, future studies should examine further and use confirmatory techniques.

Regarding to response categories, we found difficulties in distinguishing between the response options very little (1) and little (2), or much (4) and very much (5). This pattern was also reported by Munyombwe, et al.¹⁰, who suggests that it could be explained by the sample size or an excess of response categories.

According to the findings of the item-map graphics for the subscales, some patients are in the bottom of the emotional and physical subscales person-item maps, denoting floor effects. This finding is consistent with Munyombwe, et al.¹⁰, and suggests that those subscales need more items to cover all the levels of the underlying trait. Nevertheless, some studies have reported either floor or ceiling effect in the analysis of total score of the MLHF-Q¹⁰.

In relation to other variables that measure different stages of disease severity, our results are consistent with a priori hypothesis. The MLHF-Q scores clearly discriminate between different stages of NYHA functional class and age. This has also been observed in both observational studies^{9,11,13}, as well as clinical trials, where it is the ideal setting for assessing sensitivity to change³²⁻³⁴.

Strengths and limitations

The strengths of our study include an adequate sample size, as also shown by the KMO statistic. Also, we provide complete analyses of the structural validity, using both factorial PCA and Rasch analysis. The present study has, however, some important limitations to consider. First, our study was conducted in a single HF center. Accordingly, study results cannot be considered a representative description of the HRQoL of all Colombia's HF clinics. Second, the MLHF-Q is a self-administered questionnaire and, in our study it was applied by a nurse because a high percentage of our population had low educational level, and therefore it could have affected the measurement of the HRQoL.

Conclusions

In conclusion, we have assessed the content, the internal consistency, construct and discriminative capacity of the MLHF-Q in patients with CHF from Colombia. We have confirmed the three-factor structure of MLHF-Q such as previous studies, and satisfactory level for internal consistency. Additionally, these results suggest that the questionnaire adequately reflects the severity of the disease. However further studies are required in Colombian population to validate these findings and to evaluate the sensitivity to change of the MLHF-Q in longitudinal designs.

Acknowledgements

The authors are grateful to patients participating in this study.

Conflict of interest

The authors declare no conflicts of interest.

References

1. Roger VL. Epidemiology of heart failure. *Circ Res.* 2013; 113(6): 646-659. doi: 10.1161/CIRCRESAHA.113.300268.
2. Ciapponi A, Alcaraz A, Calderon M, Matta MG, Chaparro M, Soto N, et al. Burden of heart failure in Latin América: a systematic review and meta-analysis. *Rev Esp Cardiol (Engl Ed).* 2016; 69(11): 1051-1060. doi: 10.1016/j.rec.2016.04.054.
3. Escobar A, Garcia-Perez L, Navarro G, Bilbao A, Quiros R, group C-HS. A one-year mortality clinical prediction rule for patients with heart failure. *Eur J Intern Med.* 2017. doi: 10.1016/j.ejim.2017.06.013.
4. Jaarsma T, Johansson P, Agren S, Stromberg A. Quality of life and symptoms of depression in advanced heart failure patients and their partners. *Curr Opin Support Palliat Care.* 2010; 4(4): 233-237. doi: 10.1097/SPC.0b013e328340744d.
5. Juenger J, Schellberg D, Kraemer S, Haunstetter A, Zugck C, Herzog W, et al. Health related quality of life in patients with congestive heart failure: comparison with other chronic diseases and relation to functional variables. *Heart.* 2002; 87(3): 235-241.
6. Allen LA, Stevenson LW, Grady KL, Goldstein NE, Matlock DD, Arnold RM, et al. Decision making in advanced heart failure: a scientific statement from the American Heart Association. *Circulation.* 2012; 125(15): 1928-1952. doi: 10.1161/CIR.0b013e31824f2173.
7. Berry C, McMurray J. A review of quality-of-life evaluations in patients with congestive heart failure. *Pharmacoeconomics.* 1999;16(3): 247-271. doi: 10.2165/00019053-199916030-00003.
8. Rector TS, Cohn JN. Assessment of patient outcome with the Minnesota Living with Heart Failure questionnaire: reliability and validity during a randomized, double-blind, placebo-controlled trial of pimobendan. *Pimobendan Multicenter Research Group. Am Heart J.* 1992; 124(4): 1017-1025.
9. Lambrinou E, Kalogirou F, Lamnisos D, Middleton N, Sourtzi P, Lemonidou C, et al. Evaluation of the psychometric properties of the Greek version of the Minnesota Living With Heart Failure questionnaire. *J Cardiopulm Rehabil Prev.* 2013; 33(4): 229-233. doi: 10.1097/HCR.0b013e3182930cbb.
10. Munyombwe T, Hofer S, Fitzsimons D, Thompson DR, Lane D, Smith K, et al. An evaluation of the Minnesota Living with Heart Failure Questionnaire

- using Rasch analysis. *Qual Life Res.* 2014; 23(6): 1753-1765. doi: 10.1007/s11136-013-0617-0.
11. Garin O, Soriano N, Ribera A, Ferrer M, Pont A, Alonso J, et al. Validation of the Spanish version of the Minnesota Living with Heart Failure Questionnaire. *Rev Esp Cardiol.* 2008; 61(3): 251-259.
 12. Parajon T, Lupon J, Gonzalez B, Urrutia A, Altimir S, Coll R, et al. Use of the Minnesota Living With Heart Failure Quality of Life Questionnaire in Spain. *Rev Esp Cardiol.* 2004; 57(2): 155-160.
 13. Naveiro-Rilo JC, Diez-Juarez DM, Romero Blanco A, Rebollo-Gutierrez F, Rodriguez-Martinez A, Rodriguez-Garcia MA. Validation of the Minnesota living with heart failure questionnaire in primary care. *Rev Esp Cardiol.* 2010; 63(12): 1419-1427.
 14. Garin O, Herdman M, Vilagut G, Ferrer M, Ribera A, Rajmil L, et al. Assessing health-related quality of life in patients with heart failure: a systematic, standardized comparison of available measures. *Heart Fail Rev.* 2014; 19(3): 359-367. doi: 10.1007/s10741-013-9394-7.
 15. Middel B, Bouma J, de Jongste M, van Sonderen E, Niemeijer MG, Crijns H, et al. Psychometric properties of the Minnesota Living with Heart Failure Questionnaire (MLHF-Q). *Clin Rehabil.* 2001; 15(5): 489-500. doi: 10.1191/026921501680425216.
 16. Zanolta L, Zardini P. Selection of endpoints for heart failure clinical trials. *Eur J Heart Fail.* 2003; 5(6): 717-723.
 17. Garin O, Ferrer M, Pont A, Rue M, Kotzeva A, Wiklund I, et al. Disease-specific health-related quality of life questionnaires for heart failure: a systematic review with meta-analyses. *Qual Life Res.* 2009; 18(1): 71-85. doi: 10.1007/s11136-008-9416-4.
 18. Saccomann IC, Cintra FA, Gallani MC. Psychometric properties of the Minnesota Living with Heart Failure--Brazilian version--in the elderly. *Qual Life Res.* 2007; 16(6): 997-1005. doi: 10.1007/s11136-007-9170-z.
 19. Ho CC, Clochesy JM, Madigan E, Liu CC. Psychometric evaluation of the Chinese version of the Minnesota Living with Heart Failure Questionnaire. *Nurs Res.* 2007; 56(6): 441-448. doi: 10.1097/01.NNR.0000299849.21935.c4.
 20. Cronbach L. Coefficient alpha and the internal structure of test. *Psychometrika.* 1951; 16(3): 297-334.
 21. Kaiser H. An index of factorial simplicity. *Psychometrika.* 1974; 39: 32-36.
 22. Bartlett M. Test of significance in factor analysis. *Psychology.* 1950; 3: 77-85.
 23. Garin O, Ferrer M, Pont A, Wiklund I, Van Ganse E, Vilagut G, et al. Evidence on the global measurement model of the Minnesota Living with Heart Failure Questionnaire. *Qual Life Res.* 2013; 22(10): 2675-2684. doi: 10.1007/s11136-013-0383-z.
 24. Bilbao A, Escobar A, Garcia-Perez L, Navarro G, Quiros R. The Minnesota living with heart failure questionnaire: comparison of different factor structures. *Health Qual Life Outcomes.* 2016; 14: 23. doi: 10.1186/s12955-016-0425-7.
 25. Tesio L. Measuring behaviours and perceptions: Rasch analysis as a tool for rehabilitation research. *J Rehabil Med.* 2003; 35(3):105-115.
 26. Mogle J, Buck H, Zambroski C, Alvaro R, Vellone E. Cross-Validation of the Minnesota Living With Heart Failure Questionnaire. *J Nurs Scholarsh.* 2017; 49(5): 513-520. doi: 10.1111/jnu.12318.
 27. Heo S, Moser DK, Riegel B, Hall LA, Christman N. Testing the psychometric properties of the Minnesota Living with Heart Failure questionnaire. *Nurs Res.* 2005; 54(4): 265-272.
 28. Moon JR, Jung YY, Jeon ES, Choi JO, Hwang JM, Lee SC. Reliability and validity of the Korean version of the Minnesota Living with Heart Failure Questionnaire. *Heart Lung.* 2012; 41(1): 57-66. doi: 10.1016/j.hrtlng.2011.09.011.
 29. Breidhardt T, Irfan A, Klima T, Drexler B, Balmelli C, Arenja N, et al. Pathophysiology of lower extremity edema in acute heart failure revisited. *Am J Med.* 2012; 125(11): 1124 e1- e8. doi: 10.1016/j.amjmed.2011.12.015.
 30. Kataoka H. Clinical characteristics of lower-extremity edema in stage A cardiovascular disease status defined by the ACC/AHA 2001 Chronic Heart Failure Guidelines. *Clin Cardiol.* 2013; 36(9): 555-559. doi: 10.1002/clc.22159.
 31. Sztajzel J. Chronic heart failure and its consequences on the partner relationship. *Rev Med Suisse.* 2015; 11(498): 2313-2315.
 32. Kitzman DW, Hundley WG, Brubaker PH, Morgan TM, Moore JB, Stewart KP, et al. A randomized double-blind trial of enalapril in older patients with heart failure and preserved ejection fraction: effects on exercise tolerance and arterial distensibility. *Circ Heart Fail.* 2010; 3(4): 477-485. doi: 10.1161/CIRCHEARTFAILURE.109.898916.
 33. Rector TS, Carson PE, Anand IS, McMurray JJ, Zile MR, McKelvie RS, et al. Assessment of long-term effects of irbesartan on heart failure with preserved ejection fraction as measured by the minnesota living with heart failure questionnaire in the irbesartan in heart failure with preserved systolic function (I-PRESERVE) trial. *Circ Heart Fail.* 2012; 5(2): 217-225. doi: 10.1161/

CIRCHEARTFAILURE.111.964221.
 34. Edelmann F, Wachter R, Schmidt AG, Kraigher-Krainer E, Colantonio C, Kamke W, et al. Effect of spironolactone on diastolic function and exercise

capacity in patients with heart failure with preserved ejection fraction: the Aldo-DHF randomized controlled trial. JAMA. 2013;309(8): 781-791. doi: 10.1001/jama.2013.905.

Supplementary Material

Supplementary Table S1. Internal consistency of the Minnesota Living with Heart Failure Questionnaire (n=200).

Item	Sign	Item-test correlation	Item-rest correlation	Average inter-item covariance	Alpha
1. Swelling in your ankles, legs	+	0.5822	0.5341	1.0813	0.9103
2. Resting during day	+	0.7345	0.6966	1.0416	0.9066
3. Walking or climbing stairs difficult	+	0.7535	0.7141	1.0266	0.9059
4. Working around house difficult	+	0.7907	0.7561	1.0169	0.9049
5. Away from home difficult	+	0.7732	0.7353	1.0182	0.9053
6. Sleeping difficult	+	0.5716	0.5136	1.0697	0.9107
7. Relating to or doing things with friends or family difficult	+	0.6543	0.6091	1.0611	0.9086
8. Working to earn a living difficult	+	0.7733	0.7325	1.0087	0.9052
9. Recreational activities difficult	+	0.6852	0.6292	1.0242	0.9080
10. Sexual activities difficult	+	0.4893	0.4180	1.0811	0.9132
11. Eating less foods I like	+	0.4451	0.3842	1.1037	0.9133
12. Shortness of breath	+	0.6911	0.6498	1.0542	0.9078
13. Fatigue	+	0.7934	0.7627	1.0300	0.9052
14. Hospitalization	+	0.4095	0.3432	1.1080	0.9143
15. Medical costs	+	0.3486	0.2780	1.1199	0.9157
16. Side effects from medications	+	0.4084	0.3454	1.1108	0.9141
17. Feeling burden to family or friends	+	0.5497	0.4948	1.0813	0.9110
18. Feeling loss of self-control	+	0.5369	0.4816	1.0848	0.9113
19. Being worried	+	0.6079	0.5555	1.0655	0.9097
20. Difficulty concentrating or remembering	+	0.4493	0.3875	1.1017	0.9133
21. Being depressed	+	0.6208	0.5722	1.0674	0.9094
Test scale				1.0646	0.9137

Supplementary Table S2. Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy.

Item	KMO
1. Swelling in your ankles, legs	0.9469
2. Resting during day	0.9449
3. Walking or climbing stairs difficult	0.8850
4. Working around house difficult	0.9479
5. Away from home difficult	0.8805
6. Sleeping difficult	0.9451
7. Relating to or doing things with friends or family difficult	0.9522
8. Working to earn a living difficult	0.9219
9. Recreational activities difficult	0.9147
10. Sexual activities difficult	0.9334
11. Eating less foods I like	0.8967
12. Shortness of breath	0.9440
13. Fatigue	0.9132
14. Hospitalization	0.7203
15. Medical costs	0.7414
16. Side effects from medications	0.8842
17. Feeling burden to family or friends	0.8598
18. Feeling loss of self-control	0.8335
19. Being worried	0.8543
20. Difficulty concentrating or remembering	0.8377
21. Being depressed	0.8817
Overall	0.9004

KMO=Kaiser-Meyer-Olkin.

Psychometric properties of the Minnesota Living with Chronic Heart Failure Questionnaire in a Colombian population

Supplementary Table S3. Extraction of factors of the Minnesota Living with Heart Failure Questionnaire (n=200).

Dimension	Eigenvalue	Difference	Proportion	Cumulative
Physical	6.43121	3.11483	0.3062	0.3062
Emotional	3.31639	1.71791	0.1579	0.4642
Social	1.59847	.	0.0761	0.5403

Method: principal-component factors; Rotation: orthogonal varimax (Kaiser off); Likelihood Ratio test: independent vs. saturated: $\chi^2(210) = 2126.20$ Prob> $\chi^2 = 0.0000$.

Supplementary Table S4. Standardized residuals variance, eigenvalues in the 1st contrast and general coefficients of correlation between residuals (n=192).

Dimension	% Variance by measures	Eigenvalues unexplained variance in 1st contrast	Coefficients between residuals
Physical	57.4	2	All <0.50
Emotional	54.0	1.7	19-20 (-0.51)
Emotional without item 20	59.1	1.7	All <0.50
Social	44.8	1.7	All <0.50
Total	42.0	3.3	19-21 (0.52)

Supplementary Table S5. Severity levels, standard errors, and goodness-of-fit indices of the MLHF-Q emotional dimension without item 20 using Rasch analysis (n=192).


Items	Description	(logit)	SE	MNSQ	
				Infit	Outfit
17	Feeling burden to family or friends	0.29	0.08	1.21	1.18
21	Being depressed	-0.02	0.08	1.02	0.96
19	Being worried	-0.45	0.08	0.92	0.88
18	Feeling a loss of self-control	0.19	0.08	0.84	0.89

SE=Standard error; MNSQ=Mean square fit statistic; Infit=Inlier-sensitive fit; Outfit=Outlier-sensitive fit.


Supplementary Table S6. Discriminatory capacity of the Minnesota Living with Heart Failure Questionnaire (n=200).

Variable	Physical Median (Q1-Q3)	Emotional Median (Q1-Q3)	Social Median (Q1-Q3)	Total Median (Q1-Q3)
Sex				
Women	15 (4-26)	9 (4-15)	16 (9-23)	43 (23-58)
Men	10 (3-22)	7 (3-12)	16 (10-21)	35 (19-53)
p-value	0.433	0.047*	0.730	0.273
Age				
>65	18 (6-24)	9 (4-14)	17 (12-22)	45 (27-58)
≤65	8 (3-20)	6 (3-12)	15 (8-22)	29 (16-52)
p-value	0.002*	0.031*	0.182	0.005*
Functional class NYHA				
III-IV	27 (21-32)	12 (8-18)	23 (18-27)	61 (53-77)
I-II	7 (3-17)	6 (3-10)	13 (8-20)	28 (15-46)
p-value	0.000*	0.000*	0.000*	0.000*
LVEF (%)				
≤45	18 (8-23)	8 (4-14)	14 (7-20)	45 (23-58)
>45	10 (3-22)	7 (3-12)	17 (10-22)	35 (20-54)
p-value	0.146	0.267	0.177	0.556

*=p-value <0.05 Mann-Whitney U test. Q1-Q3=First and third quartile; LVEF=Left ventricular ejection fraction; NYHA=New York Heart Association.



Supplementary Figure S1. Category probability curves.



Supplementary Figure S2. Wright map for emotional, physical and social dimensions of the MLHF-Q using Rasch analysis (n=192). Each “#” = 2 people; each “.” = 1 person.